

ONTARIO CORN COMMITTEE

The ONTARIO CORN COMMITTEE is made up of representatives of Agriculture and Agri-Food Canada, the Ontario Ministry of Agriculture, Food and Rural Affairs, the University of Guelph, the Ontario Soil and Crop Improvement Association, the Grain Farmers of Ontario and the Canadian Seed Trade Association. Hybrid Performance trials are conducted each year by the following cooperating agencies:

Ridgetown Campus, University of Guelph;

Plant Agriculture Department, University of Guelph;

Winchester Research Station, University of Guelph,

Kent Ag Research Inc.,

Agriculture and Agri-Food Canada at Ottawa.

TESTING METHODS

Hybrids entered in the Hybrid Corn Performance Trials are selected by the seed companies. A testing fee is charged per hybrid per replication. A hybrid must be entered in all trials within a table.

In each trial, hybrids are replicated in a suitable experimental design. Trials are machine planted with an excess of seed and thinned at an early growth stage to obtain a uniform population. A row width of 30 inches is used in all trials. Plots consist of four rows of which the middle two rows are harvested for yield. Fertilizer rates may be higher than those recommended by OMAFRA to compensate for any variability in soil nutrient supply.

Most of the hybrids entered in the trials were treated with a seed treatment to control soil insects. Hybrids that were not treated with are not identified in the report. There was no significant damage from soil insects at any of the locations.

To determine the percentage of lodged plant, a count is made, immediately before harvest, of all plants broken below the ear and all plants which are leaning such that the ear is in the adjacent row or is otherwise unharvestable.

The moisture percentage of the grain is measured at harvest time. The weight of grain harvested from each plot is determined and the yield of shelled corn is calculated at 15% moisture. Test weights are recorded either during harvest, using combine-mounted monitoring equipment, or in the laboratory, using accepted procedures.

DUPLICATION OF THIS REPORT:

This report may be reproduced in its entirety provided that due credit is given to The Ontario Corn Committee and provided that neither the content nor the appearance of the report is altered. Tables may be reproduced provided that the entire table, with accompanying notes, is included.

For further information, contact

The Secretary, Ontario Corn Committee,
107 Templeton Court
Kemptville, ON K0G 1J0
eastcropper@gmail.com